



THE UNIVERSITY *of* EDINBURGH

## Edinburgh Research Explorer

### Prolonged ventilation in thyroid storm

**Citation for published version:**

Wake, D, Lamb, J, Babu, N & Campbell, IW 2008, 'Prolonged ventilation in thyroid storm', *Scottish Medical Journal*, vol. 53, no. 3, pp. 1-4. <https://doi.org/10.1258/RSMJMJ.53.3.571>

**Digital Object Identifier (DOI):**

[10.1258/RSMJMJ.53.3.571](https://doi.org/10.1258/RSMJMJ.53.3.571)

**Link:**

[Link to publication record in Edinburgh Research Explorer](#)

**Document Version:**

Publisher's PDF, also known as Version of record

**Published In:**

Scottish Medical Journal

**General rights**

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

**Take down policy**

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact [openaccess@ed.ac.uk](mailto:openaccess@ed.ac.uk) providing details, and we will remove access to the work immediately and investigate your claim.





## Prolonged Ventilation in Thyroid Storm

D J Wake<sup>1</sup>, J V Lamb<sup>2</sup>, N Babu<sup>3</sup>, IW Campbell<sup>2</sup>

<sup>1</sup> Metabolic Unit, Western General Hospital, Edinburgh EH4 2XU <sup>2</sup> Department of Medicine , Victoria Hospital, Kirkcaldy KY2 5AH, <sup>3</sup> Department of Anaesthetics, Victoria Hospital, Kirkcaldy KY2 5AH

Corresponding Author: Dr Deborah J Wake, Specialist Registrar, Metabolic Unit, Western General Hospital, Crewe Road, Edinburgh, EH4 2XU

Email: [debbie@thewakesplace.co.uk](mailto:debbie@thewakesplace.co.uk)

### ABSTRACT

Thyroid storm (crisis) is uncommon but may be life threatening and is recognised by an exaggeration of the clinical features of thyrotoxicosis. Proximal myopathy is a well recognised presenting feature of Graves disease. Respiratory muscle weakness may also commonly occur in thyrotoxicosis but is often undiagnosed. We report a case of thyroid storm with rapid atrial fibrillation, severe agitation and extreme widespread muscle weakness. The respiratory muscles were so compromised that a respiratory arrest occurred. Ventilation was required for 7 weeks until full recovery occurred.

Keywords: Thyroid storm, thyrotoxicosis, ventilation, respiratory failure, myopathy

## CASE HISTORY

A previously fit and well 58 year old shop worker presented as an emergency with a 3 month history of worsening breathlessness, agitation and palpitations. She was noted to have bilateral lid retraction and exophthalmos, (more so on the right than the left). She was pyrexial (38.5C), with a rapid irregular pulse of 200 beats per minute, confirmed on ECG as atrial fibrillation. She was markedly dyspnoeic at rest with arterial blood gases confirming type 1 respiratory failure (pO<sub>2</sub> -7.9kPa, pCO<sub>2</sub> -5.9kPa, O<sub>2</sub> sats of 77% on air). Blood Pressure was 140/100mmHg. The clinical impression of a thyroid storm was confirmed by a markedly elevated free thyroxine (T<sub>4</sub>; 360pmol/L (ref range 9-19)) with an undetectable TSH. Later the TSH receptor stimulating antibodies (TRABS) were elevated at 11.2 u/L (normal range 0 – 1.5u/L) confirmed Graves disease. Urea and electrolytes were normal with no evidence of hypokalaemia. She was admitted initially to the coronary care unit for control of the atrial fibrillation with intravenous propranolol, digoxin, oral carbimazole and iodine aqueous solution.

Her condition deteriorated with the first hour, with hypoventilation, hypoxia (pO<sub>2</sub>-6.9kPa, pCO<sub>2</sub>-5.4kPa, pH-7.42; O<sub>2</sub> stats 86% (on 70%O<sub>2</sub>) and hypotension (BP- 80/40mmHg). An echocardiogram could not be interpreted because of the rapid ventricular rate. The patient was markedly hypotonic, unable to support herself upright in bed and she could not lift her head to look straight ahead. She was transferred to the intensive care unit for consideration of ventilation and on route suffered a respiratory arrest with a brief episode of asystole, which responded to a 'thump' on the sternum, returning a rhythm of fast atrial fibrillation. She was intubated within 3 minutes and ventilation started.

Seven weeks of ventilation, five weeks of which required inotropic support, followed by a five week period of rehabilitation led to a full recovery. Pulmonary function tests were not performed at presentation but this non-smoker had a FEV<sub>1</sub>/FVC of 78% predicted and a total lung capacity was 4.05 litres (predicted 4.58 litres) on recovery. The echocardiogram showed normal left ventricular function at this stage. The patient was discharged on carbimazole 20mg twice daily, propranolol 40mg twice daily with atrial fibrillation controlled at 80 beats per minute. Dalteparin was used for anticoagulation whilst ventilated and warfarin commenced prior to discharge. Some 18 months later, she was well, back at work and her euthyroid status maintained on carbimazole 20mg daily.

girdle muscles and muscles of respiration may be reduced by approx 20 and 40% respectively (6). Hypoventilation may occur, associated with diaphragmatic and intercostal muscle weakness, which is reversible with treatment (7,8). We can find no other reports in the literature of prolonged ventilation being necessary for full recovery from type 1 respiratory failure associated with thyroid storm. The exact pathophysiology is unclear, but reduction in muscle mass, dysnergia of the thoracic diaphragm and changes in adrenergic muscle stimulation may contribute.

The cardiovascular system is also directly and indirectly affected by thyrotoxicosis (9), with dysrhythmias (notably atrial fibrillation) or worsening of ischaemic heart disease symptoms. Acutely, thyrotoxicosis increases cardiac output through increased contractility and improved diastolic function. Over time however left ventricular hypertrophy and cardiac failure may occur. Acute cardiomyopathies have also been described in thyrotoxic states. These may present with sudden pulmonary oedema and are reversible upon treatment of thyrotoxicosis (10,11). Milder abnormalities of left ventricular function with no cardiac symptoms have been reported which took up to six weeks to reverse following the establishment of the euthyroid state (12). The patient described had rapid atrial fibrillation and the hypotension associated with the respiratory arrest may have been due to left ventricular dysfunction, but echocardiogram was inconclusive at diagnosis, and at six weeks showed normal findings.

In conclusion, cardiac symptoms e.g. palpitations are well recognised in thyrotoxic patients. Dyspnoea occurs in about 80% of such patients and is usually attributed to cardiac causes e.g. uncontrolled atrial fibrillation, but may be due to weakness of diaphragmatic and intercostal muscles. This should be kept in mind when assessing all thyrotoxic patients, not just those presenting in extremis with hypoventilation and type 1 respiratory failure as described. If ventilation is required in such exceptional circumstances, then it may have to be prolonged. In the case described it was seven weeks before the ventilation could be safely discontinued.

The patient has given full consent for her case details and images to be used in this publication.

## References

1. Cakir M, Samanci N, Balci N, Balci MK. Musculoskeletal manifestations in patients with thyroid

5. Kung AW. Clinical review: Thyrotoxic periodic paralysis: a diagnostic challenge *J Clin Endocrinol Metab.* 2006; 91(7): 2490-5
6. Norrelund H, Hove KY, Brems-Dalgaard E, Jurik AG, Nielsen LP, Nielsen S, Jorgensen JO, Weeke J, Moller N. Muscle mass and function in thyrotoxic patients before and during medical treatment. *Clin Endocrinol (Oxf).* 1999; 51(6): 693-9.
7. Goswami R, Guleria R, Gupta AK, Gupta N, Marwaha RK, Pande JN, Kochupillai N. Prevalence of diaphragmatic muscle weakness and dyspnoea in Graves' disease and their reversibility with carbimazole therapy. *Eur J Endocrinol.* 2002; 147(3): 299-303
8. McElvaney GN, Wilcox PG, Fairbairn MS, Hilliam C, Wilkins GE, Pare PD, Pardy RL. Respiratory muscle weakness and dyspnea in thyrotoxic patients. *Am Rev Respir Dis.* 1990; 141(5 Pt 1): 1221-7.
9. Roffi M, Cattaneo F, Topol EJ. Thyrotoxicosis and the cardiovascular system; Subtle but serious effects. *Cleveland Clinical Journal of Medicine.* 2003; 70 (1); 57-63
10. Russo S, Ombriccolo E, Papadopoulou O, Paggi A. [Dilated cardiomyopathy and polymyositis. A clinical case and critical review of the literature]. *Clin Ter.* 1997; 148(1-2):57-64.
11. Goland, S. Shimon, S., Kracoff, O. Dilated Cardiomyopathy in thyrotoxicosis *Heart.* 1999; 81; 444-445
12. Forfar JC, Muir AL, Sawers SA, Toft AD. Abnormal left ventricular function in hyperthyroidism: evidence for a possible reversible cardiomyopathy. *N Engl J Med.* 1982; 307(19): 1165-70.